

AD_____

Award Number: W81XWH-04-1-0120

TITLE: The Role of I Kappa B Kinases in Prostate Carcinogenesis and the Effect of their Inhibition on Survival of Prostate Tumors

PRINCIPAL INVESTIGATOR: Michael Karin, Ph.D.

CONTRACTING ORGANIZATION: University of California, San Diego
La Jolla, CA 92093-0723

REPORT DATE: January 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01-01-2006		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 1 Jan 2005 - 31 Dec 2005	
4. TITLE AND SUBTITLE The Role of I Kappa B Kinases in Prostate Carcinogenesis and the Effect of their Inhibition on Survival of Prostate Tumors				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-04-1-0120	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Michael Karin, Ph.D. E-Mail: karinoffice@ucsd.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, San Diego La Jolla, CA 92093-0723				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT: This research project investigates the role of the IκB kinase (IKK) and NF-κB in the development of prostate cancer (CaP), and examines the possibility that IKK inhibitors can be used in CaP treatment. To reach this goal we are employing two mouse models in which either the IKKβ or the IKKα subunit of IKK are deleted or inhibited in prostate epithelial cells. We found that neither IKKβ nor IKKα are required for normal prostate development, however IKKα may play an important role in the development of advanced CaP. The study on the role of IKKβ in prostate carcinogenesis in animal models is ongoing. We also found that IKK/NF-κB activities were increased during the evolution of androgen-independent CaP, a response that could be mediated by some of androgen-regulated genes, such as TMEFF2. Furthermore, we found that a prototypical IKK inhibitor IT-3 can suppress the proliferation of human CaP cells. Although a more thorough examination of the role of IKK/NF-κB in CaP development and progression is currently underway, our results obtained during last year suggest an important role for both IKKα and IKKβ in development and progression of CaP. Therefore inhibition of either protein kinase or both would be an effective and attractive option of the treatment of CaP.					
15. SUBJECT TERMS IκB kinase (IKK), NF-κB, prostate cancer, inhibitors, carcinogenesis, treatment					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	21	19b. TELEPHONE NUMBER (include area code)

Table of Contents

Cover.....	
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Key Research Accomplishments.....	5
Reportable Outcomes.....	7
Conclusions.....	18
References.....	18

Introduction

There are two NF- κ B activation pathways. The first pathway, the classical NF- κ B activation pathway, is normally triggered in response to microbial and viral infections and exposure to proinflammatory cytokines which activate the three subunit IKK complex leading to phosphorylation-induced degradation of I κ Bs. This pathway depends mainly on the IKK β catalytic subunit. The other pathway, the alternative pathway, leads to activation of p52:RelB dimers by inducing processing of the NF- κ B2/p100 precursor protein that binds RelB in the cytoplasm. This pathway is triggered by certain members of the tumor necrosis factor (TNF) cytokine family through selective activation of IKK α homodimers by the upstream kinase NIK.

There is considerable evidence that the two IKK/NF- κ B signaling pathways are involved in carcinogenesis, cancer progression, metastasis and drug resistance. Although certain viral proteins, cancer-associated chromosomal translocations, and mutations can lead to constitutive activation of NF- κ B in cancer progenitor cells, the most common mechanism leading to NF- κ B activation during tumorigenesis depends on autocrine and paracrine production of proinflammatory cytokines. Persistent activation of NF- κ B can lead to increased production of tumor growth factors by components of the tumor stroma as well as to upregulation of anti-apoptotic genes within the cancer cell itself. This process was recently demonstrated to occur during two different mouse models of inflammation-associated cancer leading to development of colorectal cancer and hepatocellular carcinoma. We also provided evidence for a role of NF- κ B in inflammation-driven metastatic growth. In that model, as well, IKK-driven NF- κ B activation is responsible for production of growth and survival factors by stromal components (macrophages) and upregulation of anti-apoptotic genes within the cancer cell.

Prostate cancer (CaP) is one of the most common cancers in men and the second leading cause of cancer-related deaths among men in the United States. It was shown that NF- κ B transcription factors can directly interact with several members of the nuclear receptor family including androgen receptor (AR) itself. Thus, NF- κ B may function as a co-activator for AR causing it to be active independently of androgen binding. In this case, prostate epithelial cells with high NF- κ B activity are rendered resistant to androgen withdrawal. However, NF- κ B can also trans-repress ligand-bound AR or repress the AR gene itself. Most importantly, NF- κ B activity itself is repressed by androgen treatment via AR-mediated trans-repression or other mechanisms. In this case, NF- κ B activity may increase in response to androgen withdrawal. Once activated, NF- κ B can stimulate production of various cytokines by prostate epithelial cells, CaP and stromal components. Specifically, IL-6, a well-known autocrine and paracrine growth factor for AI CaP cells, is encoded by a typical NF- κ B target gene. The dependence of IL-6 expression on NF- κ B was demonstrated in the prostate epithelium. Importantly, the human AD CaP cell line, LNCaP, was shown to assume a neuroendocrine (NE) phenotype in response to IL-6 exposure. Although NE cancers of the prostate are rare, foci with NE-like features can be observed in nearly all prostate adenocarcinomas and extensive NE differentiation is generally considered to be of poor prognostic value. Indeed, NE differentiation appears more frequently in hormone-refractory cancer. Inhibition of IL-6 activity induces the regression of AI human CaP xenografts in mice. Thus, NF- κ B inhibition should result in a similar effect by inhibiting IL-6 expression. Studies performed on various CaP cell lines reveal that AI cells often display constitutive NF- κ B activity. Furthermore, constitutively activated NF- κ B in three different human CaP cell lines was linked to overexpression of IKK subunits, and inhibition of NF- κ B activity in these cells through expression of a non-degradable super-repressor mutant of I κ B α resulted in either spontaneous apoptosis or increased sensitivity to TNF α .

SPECIFIC AIMS

- Construct mice with a conditionally activated Ikk β allele and use them to determine whether constitutive NF- κ B activation in prostate epithelial cells promotes prostate cancer.

- Construct mice with a specific deletion of *Ikkβ* in prostate epithelial cells and examine whether this deletion, as well as the inactivation of *IKKα*, inhibits prostate carcinogenesis in the TRAMP model.
- Use various strategies to inhibit *IKK* activity and test them for their ability to inhibit proliferation and induce apoptosis in prostate cancer cell lines of human origin.

Key Research Accomplishments

1. Examine whether inactivation of *IKKα* inhibits prostate carcinogenesis in the TRAMP model.

To investigate the role of the *IKKα* subunit, which may offer a more attractive target for drug development than *IKKβ* as it is not required for innate immune responses, TRAMP mice were crossed with *Ikkα^{AA/AA}* mice which express a form of *IKKα* that can not be activated because the two serines in its activation loop, which are phosphorylated by the upstream activating kinase NIK, were replaced with alanines. The resultant *Ikkα^{AA/AA}/TRAMP* mice were monitored for tumor development and found to exhibit fewer metastases (including metastasis to lymph nodes and other organs) and survive considerably longer than *WT/TRAMP* mice (Fig. 1 and Table 1). The CaP in *Ikkα^{AA/AA}/TRAMP* mice exhibit reduced cell proliferation (Fig. 2). However, there are no differences in size and weight of the prostate glands between 12-week-old *Ikkα^{AA/AA}/TRAMP* and *WT/TRAMP* mice, and both strains developed prostate adenocarcinomas, suggesting that *IKKα* kinase activity is required for CaP progression but not for normal prostate development and early tumorigenesis (Fig. 2).

We also found that the expression of some cyclins (cyclin A, cyclin B) was decreased in *Ikkα^{AA/AA}/TRAMP* tumors as compared with *WT/TRAMP* tumors (Fig. 3). A metastasis suppressor Maspin was down-regulated in wt tumors but not in *Ikkα^{AA/AA}* tumors (Fig. 4), suggesting that *IKKα* activation might repress Maspin expression which was supported by *in vitro* Maspin-luciferase reporter experiments (Fig. 5).

2. Construct mice with a specific deletion of *Ikkβ* in prostate epithelial cells and examine whether this deletion inhibits prostate carcinogenesis in the TRAMP model, and whether *IKKβ* play a role in the transition from AD CaP to AI CaP

To test the role of the *IKKβ* subunit in CaP development, we constructed a mouse strain that contains a prostate epithelium-specific deletion of the gene coding for the *IKKβ* catalytic subunit. In these experiments we took advantage of the *Ikkβ^{F/F}* mouse strain, which harbors a conditional loss-of-function “floxed” *Ikkβ* allele. To delete *IKKβ* in prostate epithelial cells we crossed *Ikkβ^{F/F}* mice to *PB-CRE4* transgenic mice, which express CRE recombinase in prostate epithelial cells. This yielded an *Ikkβ^{F/+}PB-CRE4* heterozygote strain, which after intercrossing generated a homozygote *Ikkβ^{F/F}PB-CRE4* mouse. We examined the efficiency of *IKKβ* deletion in the prostate and in purified prostate epithelial cells from 10-12 week old male *Ikkβ^{F/F}/PB-CRE4* mice by polymerase chain reaction (PCR) and immunoblotting and found efficient deletion of the *Ikkβ^F* allele and absence of *IKKβ* protein in purified ventral and dorsolateral prostate gland epithelial cells from *Ikkβ^{F/F}/PB-CRE4* mice (Fig. 6). No differences in the size of the prostate gland and its histological composition between *Ikkβ^{F/F}* and *Ikkβ^{F/F}/PB-CRE4* mice were observed. Thus, *IKKβ* is not required for normal prostate development and maintenance. Since effective and prostate-specific deletion of *IKKβ* has been confirmed, we crossed recombinant *Ikkβ^{F/F}PB-CRE4* mice as well as *Ikkβ^{F/F}* mice with the TRAMP transgenic mouse to generate *Ikkβ^{F/F}PB-CRE4-TRAMP* and *Ikkβ^{F/F}-TRAMP* mice. Cohorts of 15 male mice of the appropriate genotypes (*Ikkβ^{F/F}PB-CRE4-TRAMP* and *Ikkβ^{F/F}-TRAMP*) are currently being monitored for external signs of prostate cancer formation. Mice that will exhibit large palpable tumors will be sacrificed and both primary and metastatic tumor tissues will be collected for primary cell culture, histological and biochemical analyses.

To examine whether *IKK/NF-κB* is activated during the evolution of AI CaP, the androgen-dependent

(AD) human CaP cell line LNCaP was inoculated subcutaneously into immunocompromised SCID mice. When tumor mass reached 1 cm³, one half of the tumor-bearing mouse cohort was sacrificed and AD tumor tissue was collected. The other half of the mouse cohort was castrated. Two months later when the tumors in the castrated mice re-grew, mice were sacrificed and tumor tissues were collected to yield AI tumors. Protein lysates derived from AD and AI tumors were assayed for NF- κ B DNA binding activity by an electrophoretic mobility shift assay (EMSA) and for IKK activity by an immunocomplex kinase assay. The results clearly demonstrate that both NF- κ B and IKK activities are markedly elevated in AI tumor tissues (Fig. 7).

These results support the notion that the IKK-NF- κ B pathway may play an important role in development of AI CaP. It was reported that 80% of TRAMP mice castrated at 12 weeks of age will develop aggressive AI CaP at 24 weeks of age. To determine the effect of IKK β ablation on development of androgen-independent prostate cancer (AI CaP) in TRAMP mice, cohorts of 20 male mice of each genotype (*Ikk β ^{F/F}/PB-CRE4-TRAMP* and *Ikk β ^{F/F}/TRAMP*) are being prepared and will be castrated at 12 weeks of age and sacrificed at 24 weeks of age. Size, weight and histology of the prostate tissue and of primary and metastatic CaPs will be measured and recorded. Prostate tissues, primary and metastatic CaP will be collected for primary cell culture, histological and biochemical analyses.

While we have been breeding the mice described above, we have cultured primary mouse CaP cells isolated from *Ikk β ^{F/F}/TRAMP* mice, which contain two “floxed” *Ikk β* alleles that can be deleted by expression of CRE-recombinase. These cells are mixtures of tumor stroma cells and epithelial cancer cells. Cells were infected with CRE-adenovirus to delete both *Ikk β* alleles in both cell types or infected with GFP-adenovirus as a control. IKK β deletion efficiency was examined by Western blotting. *Ikk β* -deleted cells and non-deleted cells were transplanted into the flank of male *Rag1*^{-/-} mice. Two months later mice were sacrificed and tumor weights were measured. Some of the mice were castrated one month after transplantation, and two months after castration mice were sacrificed and tumor weights were measured. We found that *Ikk β* deletion in tumor cells decreased tumor growth after castration (Fig. 8).

The stroma and extracellular matrix are essential for functional and morphological differentiation of the prostatic epithelium. It is also postulated that the prostate stroma may play an important role in CaP development, as we found that the ratio between the stroma and epithelial cancer cells was increased in castrated TRAMP prostate tumors (Fig. 9). We found that both IKK and NF- κ B activities were increased in castrated CaP (Fig. 10), and *Ikk β* deletion in tumor tissue cells (both stroma and epithelial cancer cells) decreased tumor growth after castration. Analysis of the role of IKK β either in prostate epithelial cancer cells or stromal components in AI CaP development will provide us with important information as to the cell-type specificity of IKK β function. We also separated epithelial cancer cells and tumor stromal cells from CaP isolated from *Ikk β ^{F/F}/TRAMP* mice. IKK β was deleted by infection with CRE-adenovirus in either stromal cells or in cancer cells. Equal number of epithelial cancer cells and stromal cells of different genotypes will be mixed in different combinations (*Ikk β ^{F/F}* stroma+ *Ikk β ^{-/-}* CaP; *Ikk β ^{-/-}* stroma + *Ikk β ^{F/F}* CaP; etc). To examine the tumor development, equal numbers of cell combinations will be implanted under the renal capsule of male SCID mice (5 mice for each cell combination). One month later, mice will be castrated, two months later, mice will be sacrificed, and the size and weight of each tumor will be measured and recorded. Tumor tissues will be collected and analyzed. These experiments are ongoing.

**Table 1: Incidence of metastasis at death of WT/TRAMP
and IKK α AA/TRAMP mice**

	PLA(%)	Kidney+nodes (%)	Liver (%)	Lung (%)
WT/TRAMP (n=23)	20 (87)	10 (43)	4 (17)	5 (22)
IKKαAA/TRAMP (n=22)	9 (41)	4 (18)	0 (0)	0 (0)

Fig. 1





Fig. 2







Fig. 3

A



P = Primary tumor;
L = Lympho node metastasis;
Li = Liver metastasis

B



C





Fig. 4



Maspin immunostaining




Fig. 5






Fig 6, IKK β protein is efficiently deleted in prostate epithelial cells in IKK $\beta^{F/F}$ PB-CRE4 mice. Purified ventral and dorsolateral prostate epithelial cells from 10-12 week old male IKK $\beta^{F/F}$ and IKK $\beta^{F/F}$ PB-CRE4 mice were lysed for immunoblot analysis. The results show efficient deletion of IKK β in ventral and dorsolateral prostate epithelial cells from IKK $\beta^{F/F}$ PB-CRE4 mice .






Fig 7. Human LNCaP cells were implanted subcutaneously into SCID mice. When the tumor mass reached 1 cm³, half of the tumor-bearing mice were sacrificed and tumor tissues were collected (androgen-dependent tumors, AD). The remaining half were castrated. Two months later when the tumors in the castrated mice regrew, the mice were sacrificed and tumors were collected (androgen-independent tumors, AI). Tumor extracts were used for electrophoretic mobility shift assay (EMSA) (a) and determining of IKK kinase activity (b). Both NF- κ B DNA binding activity and IKK kinase activity were markedly increased in AI tumors as compared to AD tumors.

Figure 8

Deletion of IKK β in primary mouse tumor tissue cells inhibits tumor growth when host Rag1 $^{-/-}$ mice are castrated




Primary tumor tissue cells were inoculated in the flank of Rag1 $^{-/-}$ mice, two months later mice were sacrificed and tumor weight was measured




Primary tumor tissue cells were inoculated in the flank of Rag1 $^{-/-}$ mice, one month later mice were castrated. After two months of castration mice were sacrificed and tumor weight was measured

Figure 9


Pathology of prostate adenocarcinoma in castrated TRAMP mice




Primary prostate adenocarcinoma, 5x




Primary prostate adenocarcinoma, 20x



prostate adenocarcinoma, lung metastasis 20x




prostate adenocarcinoma, liver metastasis 20x



prostate adenocarcinoma, lymph node metastasis 20x


Figure 10

A



EMSA using tissue lysates from normal mouse prostate or mouse prostate cancer
P = Primary tumor;
L = Lympho node metastasis;
LI = Liver metastasis

B



IKK Kinase Assay using tissue lysates from normal mouse prostate or mouse prostate cancer
P = Primary tumor;
L = Lympho node metastasis;
LI = Liver metastasis

Conclusions

In summary, results obtained during the past project year suggest that both IKK α and IKK β may play an important role in development and progression of CaP. Clearly, a more thorough examination of the role of the two IKK subunits in CaP development and progression is needed, especially during the transition of CaP from an AD to an AI state. These are functions being examined during the current project year. Most interesting is the finding of maspin upregulation in CaP from *Ikk $\alpha^{aa/aa}$ /TRAMP* mice. If maspin will be verified to be a tumor and metastases suppressor in this model as well as in human CaP, our work provides a clear outline for the development of new therapeutic approach to prostate cancer.

REFERENCES

1. Ghosh, S., May, M.J. & Kopp, E.B. NF-kB and Rel proteins: evolutionarily conserved mediators of immune responses. *Ann Rev Immunol* 16, 225-260 (1998).
2. Gilmore, T.D. Multiple mutations contribute to the oncogenicity of the retroviral oncoprotein v-Rel. *Oncogene* 18, 6925-6937 (1999).
3. Karin, M. & Ben-Neriah, Y. Phosphorylation meets ubiquitination: the control of NF-kB activity. *Annu Rev Immunol* 18, 621-663 (2000).
4. Li, Z.-W. et al. The IKK β subunit of I κ B kinase (IKK) is essential for NF-kB activation and prevention of apoptosis. *J. Exp. Med.* 189, 1839-1845 (1999b).
5. Solan, N.J., Miyoshi, H., Bren, G.D. & Paya, C.V. RelB cellular regulation and transcriptional activity are regulated by p100. *J Biol Chem* 277, 1405-1418 (2002).
6. Senftleben, U. et al. Activation by IKK α of a second, evolutionary conserved, NF-kB signaling pathway. *Science* 293, 1495-1499 (2001).
7. Ghosh, S. & Karin, M. Missing pieces in the NF-kB puzzle. *Cell* 109, S81-96 (2002).
8. Karin, M. & Lin, A. NF-kB at the crossroads of life and death. *Nat Immunol* 3, 221-227 (2002).
9. Hanahan, D. & Weinberg, R.A. The hallmarks of cancer. *Cell* 100, 57-70 (2000).
10. Karin, M., Cao, Y., Greten, F. & Li, Z. NF-kB in Cancer: From innocent bystander to major culprit. *Nat Rev Cancer* 2, 301-310 (2002).
11. Xiao, G. et al. Retroviral oncoprotein Tax induces processing of NF-kB2/p100 in T cells: evidence for the involvement of IKK α . *Oncogene* 20, 6805-6815 (2001).
12. Mosialos, G. The role of Rel/NF-kB proteins in viral oncogenesis and the regulation of viral transcription. *Semin Cancer Biol* 8, 121-129 (1997).
13. Guttridge, D.C., Albanese, C., Reuther, J.Y., Pestell, R.G. & Baldwin Jr., A.S. NF-kB controls cell growth and differentiation through transcriptional regulation of cyclin D1. *Mol Cell Biol* 19, 5785-5799 (1999).
14. Hinz, M. et al. NF-kB function in growth control: regulation of cyclin D1 expression and G0/G1-to-S-phase transition. *Mol Cell Biol* 19, 2690-2698 (1999).
15. Cao, Y. et al. IKK α provides an essential link between RANK signaling and cyclin D1 expression during mammary gland development. *Cell* 107, 763-775 (2001).
16. Beg, A.A. & Baltimore, D. An essential role for NF-kB in preventing TNF- α induced cell death. *Science* 274, 782-784 (1996).

17. Liu, Z.-G., Hu, H., Goeddel, D.V. & Karin, M. Dissection of TNF receptor 1 effector functions: JNK activation is not linked to apoptosis, while NF- κ B activation prevents cell death. *Cell* 87, 565-576 (1996).
18. Van Antwerp, D.J., Martin, S.J., Kafri, T., Green, D.R. & Verma, I.M. Suppression of TNF α -induced apoptosis by NF- κ B. *Science* 274, 787-789 (1996).
19. Wang, C.-Y., Mayo, M.W. & Baldwin, A.S., Jr. TNF- and cancer therapy-induced apoptosis: potentiation by inhibition of NF- κ B. *Science* 274, 784-787 (1996).
20. Wang, C.Y., Cusack, J.C., Jr., Liu, R. & Baldwin, A.S., Jr. Control of inducible chemoresistance: enhanced anti-tumor therapy through increased apoptosis by inhibition of NF- κ B. *Nat Med* 5, 412-417 (1999).
21. Levine, A.J. p53, the cellular gatekeeper for growth and division. *Cell* 88, 323-331 (1997).
22. Webster, G.A. & Perkins, N.D. Transcriptional cross talk between NF- κ B and p53. *Mol Cell Biol* 19, 3485-3495 (1999).
23. Tergaonkar, V., Pando, M., Vafa, O., Wahl, G. & Verma, I. p53 stabilization is decreased upon NF- κ B activation: a role for NF- κ B in acquisition of resistance to chemotherapy. *Cancer Cell* 1, 493-503 (2002).
24. Koch, A.E. et al. Interleukin-8 as a macrophage-derived mediator of angiogenesis. *Science* 258, 1798-1801 (1992).
25. Takeshita, H. et al. Matrix metalloproteinase 9 expression is induced by Epstein-Barr virus latent membrane protein 1 C-terminal activation regions 1 and 2. *J Virol* 73, 5548-5555 (1999).
26. Wang, W., Abbruzzese, J.L., Evans, D.B. & Chiao, P.J. Overexpression of urokinase-type plasminogen activator in pancreatic adenocarcinoma is regulated by constitutively activated RelA. *Oncogene* 18, 4554-4563 (1999).
27. Bond, M., Fabunmi, R.P., Baker, A.H. & Newby, A.C. Synergistic upregulation of metalloproteinase-9 by growth factors and inflammatory cytokines: an absolute requirement for transcription factor NF- κ B. *FEBS Lett* 435, 29-34 (1998).
28. Huang, S., Robinson, J.B., Deguzman, A., Bucana, C.D. & Fidler, I.J. Blockade of NF- κ B signaling inhibits angiogenesis and tumorigenicity of human ovarian cancer cells by suppressing expression of vascular endothelial growth factor and interleukin 8. *Cancer Res* 60, 5334-5339 (2000).
29. Grossmann, M.E., Huang, H. & Tindall, D.J. Androgen receptor signaling in androgen-refractory prostate cancer. *J Natl Cancer Inst* 93, 1687-1697 (2001).
30. Palvimo, J.J. et al. Mutual transcriptional interference between RelA and androgen receptor. *J Biol Chem* 271, 24151-24156 (1996).
31. Supakar, P.C., Jung, M.H., Song, C.S., Chatterjee, B. & Roy, A.K. NF- κ B functions as a negative regulator for the rat androgen receptor gene and NF- κ B activity increases during the age-dependent desensitization of the liver. *J Biol Chem* 270, 837-842 (1995).
32. McKay, L.I. & Cidlowski, J.A. Cross-talk between NF- κ B and the steroid hormone receptors: mechanisms of mutual antagonism. *Mol Endocrinol* 12, 45-56 (1998).
33. Keller, E.T., Chang, C. & Ersler, W.B. Inhibition of NF κ B activity through maintenance of I κ B α levels contributes to dihydrotestosterone-mediated repression of the interleukin-6 promoter. *J Biol Chem* 271, 26267-26275 (1996).
34. Smith, P.C. & Keller, E. Anti-interleukin-6 monoclonal antibody induces regression of human prostate cancer xenografts in nude mice. *Prostate* 48, 47-53 (2001).

35. Giri, D., Ozen, M. & Ittmann, M. Interleukin-6 is an autocrine growth factor in human prostate cancer. *Am J Pathol* 159, 2159-2165 (2001).
36. Barnes, P.J. & Karin, M. NF- κ B - A pivotal transcription factor in chronic inflammatory diseases. *New Engl. J. Med.* 336, 1066-1071 (1997).
37. Nishimura, K. et al. Potential mechanism for the effects of dexamethasone on growth of androgen-independent prostate cancer. *J Natl Cancer Inst* 93, 1739-1746 (2001).
38. Palayoor, S.T., Youmell, M.Y., Calderwood, S.K., Coleman, C.N. & Price, B.D. Constitutive activation of I κ B kinase alpha and NF- κ B in prostate cancer cells is inhibited by ibuprofen. *Oncogene* 18, 7389-7394 (1999).
39. Gasparian, A.V. et al. The role of IKK in constitutive activation of NF- κ B transcription factor in prostate carcinoma cells. *J Cell Sci* 115, 141-151 (2002).
40. Li, Q., Van Antwerp, D., Mercurio, F., Lee, K.-F. & Verma, I.M. Severe liver degeneration in mice lacking the I κ B kinase 2 gene. *Science* 284, 321-325 (1999a).
41. Senftleben, U., Li, Z.-W., Baud, V. & Karin, M. IKK β is essential for protecting T cells from TNF α -induced apoptosis. *Immunity* 14, 217-230 (2001).
42. Makris, C. et al. Female mice heterozygote for IKK γ /NEMO deficiencies develop a genodermatosis similar to the human X-linked disorder Incontinentia Pigmenti. *Mol. Cell* 15, 969-979 (2000).
43. Park, J.M., Greten, F.R., Li, Z.-W. & Karin, M. Macrophage apoptosis by Anthrax Lethal Factor and p38 MAP kinase inhibition. *Science* 297, 2048-2051 (2002).
44. Chen, L.-W. et al. The two faces of IKK and NF- κ B inhibition: Prevention of systemic inflammation but increased local injury following intestinal ischemia-reperfusion. *Nat Med* In Press(2003).
45. Bargou, R.C. et al. Constitutive NF- κ B-RelA activation is required for proliferation and survival of Hodgkin's disease tumor cells. *J. Clin. Invest.* 100, 2961-2969 (1997).
46. Cahir-McFarland, E.D., Davidson, D.M., Schauer, S.L., Duong, J. & Kieff, E. NF- κ B inhibition causes spontaneous apoptosis in Epstein-Barr virus-transformed lymphoblastoid cells. *Proc Natl Acad Sci U S A* 97, 6055-6060 (2000).
47. Keller, S.A., Schattner, E.J. & Cesarman, E. Inhibition of NF- κ B induces apoptosis of KSHV-infected primary effusion lymphoma cells. *Blood* 96, 2537-2542 (2000).
48. Furman, R.R., Asgary, Z., Mascarenhas, J.O., Liou, H.C. & Schattner, E.J. Modulation of NF- κ B activity and apoptosis in chronic lymphocytic leukemia B cells. *J Immunol* 164, 2200-2206 (2000).
49. Baldwin, A.S. Control of oncogenesis and cancer therapy resistance by the transcription factor NF- κ B. *J Clin Invest* 107, 241-246 (2001).
50. Valen, G., Yan, Z.Q. & Hansson, G.K. NF- κ B and the heart. *J Am Coll Cardiol* 38, 307-314 (2001).
51. Haefner, B. NF- κ B: arresting a major culprit in cancer. *Drug Discov Today* 7, 653-663 (2002).
52. May, M.J. et al. Selective inhibition of NF- κ B activation by a peptide that blocks the interaction of NEMO with the I κ B kinase complex. *Science* 289, 1550-1554 (2000).
53. Wu, X. et al. Generation of a prostate epithelial cell-specific Cre transgenic mouse model for tissue-specific gene ablation. *Mech Dev* 101, 61-69 (2001).

54. Hu, Y. et al. Abnormal morphogenesis but intact IKK activation in mice lacking the IKK α subunit of the I κ B kinase. *Science* 284, 316-320 (1999).
55. Berges, R.R. et al. Cell proliferation, DNA repair, and p53 function are not required for programmed death of prostatic glandular cells induced by androgen ablation. *Proc Natl Acad Sci U S A* 90, 8910-8914 (1993).
56. Buttyan, R., Ghafar, M.A. & Shabsigh, A. The effects of androgen deprivation on the prostate gland: cell death mediated by vascular regression. *Curr Opin Urol* 10, 415-420 (2000).
57. Greenberg, N.M. et al. Prostate cancer in a transgenic mouse. *Proc Natl Acad Sci U S A* 92, 3439-3443 (1995).
58. Li, Z.-W., Omori, S.A., Labuda, T., Karin, M. & Rickert, R.C. IKK β is required for peripheral B cell proliferation and survival. *J Immunol* 170, In Press (2003).
59. Callahan, J.E. & Roshak, A.K. NF- κ B inhibitors. in *Int. Pat. Appl. PCT Gazette WO 02/30353 A2* (2002).
60. Hu, Y. et al. IKK α controls formation of the epidermis independently of NF- κ B via a differentiation inducing factor. *Nature* 410, 710-714 (2001).
61. Wender, P.A. et al. The design, synthesis, and evaluation of molecules that enable or enhance cellular uptake: peptoid molecular transporters. *Proc Natl Acad Sci U S A* 97, 13003-13008 (2000).
62. Rothbard, J.B. et al. Arginine-rich molecular transporters for drug delivery: role of backbone spacing in cellular uptake. *J Med Chem* 45(2002).
63. DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. & Karin, M. A cytokine-responsive I κ B kinase that activates the transcription factor NF- κ B. *Nature* 388, 548-554 (1997).
64. Zandi, E., Rothwarf, D.M., Delhase, M., Hayakawa, M. & Karin, M. The I κ B kinase complex (IKK) contains two kinase subunits, IKK α and IKK β , necessary for I κ B phosphorylation and NF- κ B activation. *Cell* 91, 243-252 (1997).
65. Delhase, M., Hayakawa, M., Chen, Y. & Karin, M. Positive and negative regulation of I κ B kinase activity through IKK β subunit phosphorylation. *Science* 284, 309-313 (1999).
66. Yeh, R.H., Lee, T.R. & Lawrence, D.S. From consensus sequence peptide to high affinity ligand, a "library scan" strategy. *J Biol Chem* 276, 12235-12240 (2001).
67. Pinilla, C., Appel, J.R., Borrás, E. & Houghten, R.A. Advances in the use of synthetic combinatorial chemistry: Mixture-based libraries. *Nat Med* 9, 118-122 (2003).
68. Lee, P.H. & Bevis, D.J. Development of a homogeneous high throughput fluorescence polarization assay for G protein-coupled receptor binding. *J Biomol Screen* 5, 415-419 (2000).
69. Kallunki, T., Deng, T., Hibi, M. & Karin, M. c-Jun can recruit JNK to phosphorylate dimerization partners via specific docking interactions. *Cell* 87, 929-939 (1996).
70. Songyang, Z. et al. Use of an oriented peptide library to determine the optimal substrates of protein kinases. *Current Biology* 4, 973-982 (1994).